

EXPLICIT POINTS ON THE LEGENDRE CURVE

DOUGLAS ULMER

ABSTRACT. Using explicit points and elementary arguments we exhibit non-isotrivial elliptic curves over function fields with Mordell-Weil groups of arbitrarily large rank.

1. INTRODUCTION

There are now several constructions of elliptic curves (and higher-dimensional Jacobians) of large rank over $\mathbb{F}_p(t)$ or $\overline{\mathbb{F}}_p(t)$. Most of these constructions rely on relatively sophisticated mathematics, such as the theory of algebraic surfaces, cohomology, and L -functions. See [UlmPCMI] for an overview of some of these techniques. See also [Con09] where Conceição gives an explicit construction of polynomial points on quadratic twists of constant supersingular elliptic curves over $\mathbb{F}_p(t)$ and uses them to show that the rank is large.

Our aim in this short note is to give an elementary and explicit construction of many independent points on non-isotrivial elliptic curves over $\mathbb{F}_p(t)$ ($p > 2$) with arbitrarily large rank. More precisely, we consider the Legendre curve

$$(1.1) \quad y^2 = x(x-1)(x-t)$$

over $\mathbb{F}_p(t)$ and over extensions $\mathbb{F}_q(t^{1/d})$. That one should expect this curve to have unbounded ranks in these towers follows from [Ulm07, 4.7] and the Birch and Swinnerton-Dyer conjecture, but so far the Legendre curve does not seem to fall into the class of curves constructed in [Ber08] and studied in [UlmDPCT]. It came as something of a surprise to the author that one can nevertheless write down explicit points in the simplest way imaginable.

This note came about as a result of conversations with Dick Gross at the 2009 IAS/Park City Mathematics Institute. It is a pleasure to thank the organizers of the meeting for the opportunity to give a course, the audience for their attention and comments, and Gross for his interest and incisive questions. Thanks are also due to Chris Hall for thought-provoking data on L -functions and stimulating conversations on questions around this paper.

2. POINTS

Throughout the paper, p will be an odd prime number and E will be the elliptic curve

$$(2.1) \quad E: \quad y^2 = x(x+1)(x+t)$$

Date: December 14, 2009.

This paper is based upon work partially supported by the National Science Foundation under Grant No. DMS 0701053.

over $\mathbb{F}_p(t)$ or one of its extensions. Over any extension with a primitive 4th root of unity ζ_4 , E is isomorphic to the Legendre curve (1.1) via the change of coordinates $(x, y) \mapsto (-x, \zeta_4 y)$.

The j invariant of E is

$$j(E) = \frac{256(t^2 - t + 1)^3}{t^2(t - 1)^2}$$

and the discriminant of the model (2.1) is

$$\Delta = 16t^2(t - 1)^2.$$

It is well-known, and we will reprove below, that over $\overline{\mathbb{F}}_p(t)$, E has exactly four points: the point at infinity and the 2-torsion points $(0, 0)$, $(-1, 0)$, and $(-t, 0)$.

We will consider E over various extensions $\mathbb{F}_q(t^{1/d}) = \mathbb{F}_q(u)$ of $\mathbb{F}_p(t)$ with $u^d = t$. The most important case will be when d has the form $d = p^f + 1$ and the d -th roots of unity μ_d lie in $\mathbb{F}_q^\times$. (Note that if $d = p^f + 1$, $f \geq 1$, then $\mathbb{F}_p(\mu_d) = \mathbb{F}_{p^{2f}}$.) We write K_d for the extension $\mathbb{F}_p(\mu_d, u)$ of $\mathbb{F}_p(t)$ with $u^d = t$. We fix a primitive d -th root of unity in K_d and denote it ζ_d .

Over K_d with d of the form $d = p^f + 1$, $f \geq 0$, E has the Weierstrass equation

$$y^2 = x(x + 1)(x + u^{p^f + 1})$$

and the obvious point $P(u) = (u, u(u + 1)^{(p^f + 1)/2})$. Since E is defined over $\mathbb{F}_p(t) \subset \mathbb{F}_p(\mu_d, t)$, applying elements of $\text{Gal}(K_d/\mathbb{F}_p(\mu_d, t)) \cong \mu_d$ to $P(u)$ we get a collection of points

$$(2.2) \quad P_i = P(\zeta_d^i u) = (\zeta_d^i u, \zeta_d^i u(\zeta_d^i u + 1)^{(p^f + 1)/2}) \quad i \in \mathbb{Z}/d\mathbb{Z}.$$

We will see below by elementary means that these points are almost independent—they generate a subgroup of $E(K_d)$ of rank $d - 2 = p^f - 1$. Using somewhat more technology, we'll see in the following sections that they generate a subgroup of finite index, i.e., that the rank of $E(K_d)$ is $d - 2$.

3. TORSION

We briefly detour to find all torsion points on $E(\overline{\mathbb{F}}_p(t^{1/d}))$ for all d prime to p .

It is clear that $E(\mathbb{F}_p(t))$ has 3 points of order exactly 2. Let us label the points $Q_0 = (0, 0)$, $Q_1 = (-1, 0)$, and $Q_t = (-t, 0)$. It is also easy to see that $E(K_2)$ has 4 points of order 4, namely $\pm P_0$ and $\pm P_1$ in the notation of (2.2). We have

$$2P_0 = Q_0 \quad P_0 + Q_0 = -P_0 \quad P_0 + Q_1 = P_1 \quad P_0 + Q_t = -P_1.$$

These turn out to be all the torsion points rational in the tower $\overline{\mathbb{F}}_p(t^{1/d})$:

3.1. Proposition. *Let d be a positive integer not divisible by p . If d is odd, then $E(\overline{\mathbb{F}}_p(t^{1/d}))_{\text{tor}} \cong (\mathbb{Z}/2\mathbb{Z}) \times (\mathbb{Z}/2\mathbb{Z})$. If d is even, then $E(\overline{\mathbb{F}}_p(t^{1/d}))_{\text{tor}} \cong (\mathbb{Z}/2\mathbb{Z}) \times (\mathbb{Z}/4\mathbb{Z})$.*

Proof. We give a brief sketch of the proof. The prime-to- p part of the proposition also follows from the Birch and Swinnerton-Dyer formula and the height computation below.

We have already exhibited points showing that the torsion subgroup is at least as large as claimed.

Since the j invariant of E is not a p -th power in $\overline{\mathbb{F}}_p(t^{1/d})$ for any d prime to p , E can have no p -torsion over these fields.

If d is odd, we will see below that E has reduction type I_{2d}^* at $u = \infty$. Thus the prime-to- p torsion must be a subgroup of $(\mathbb{Z}/2\mathbb{Z}) \times (\mathbb{Z}/2\mathbb{Z})$ and so is equal to this group.

This argument does not work for d even, since the reduction at ∞ is of type I_{2d} . For d even or odd, we may argue as follows: If $E(\overline{\mathbb{F}}_p(t^{1/d}))$ had a point of order ℓ for some $\ell \nmid 2p$, we would get a non-constant map from the $u = t^{1/d}$ -line to a modular curve of positive genus; this is clearly impossible. To rule out any additional 2-power torsion, one may use the duplication formula to see that the solutions R of $2R = P_0$, $2R = P_1$, and $2R = Q_1$ are defined over fields which are ramified over a finite non-zero place of $\overline{\mathbb{F}}_p(t^{1/2})$. Since $\overline{\mathbb{F}}_p(t^{1/d})$ is unramified over $\overline{\mathbb{F}}_p(t^{1/2})$ away from 0 and ∞ , no such R is rational over $\overline{\mathbb{F}}_p(t^{1/d})$.

This completes the sketch of proof of the proposition. $\square$

4. HEIGHTS

The height pairing on $E(K_d)$ is a symmetric bilinear form which is non-degenerate modulo torsion. It is closely related to the intersection pairing on the surface $\mathcal{E} \rightarrow \mathbb{P}^1$ associated to E . In this section we will use it to find the relations among the points P_i modulo torsion.

4.1. The Néron model. Fix an integer d , let $K_d = \mathbb{F}_p(\mu_d, u)$, and consider E over K_d . In order to compute heights on $E(K_d)$ we need a nice model for E over $\mathbb{P}^1$, specifically, a smooth proper surface $\mathcal{E}$ over $\mathbb{F}_q = \mathbb{F}_p(\mu_d)$ equipped with a relatively minimal morphism $\pi : \mathcal{E} \rightarrow \mathbb{P}^1$ whose generic fiber is E .

The construction of $\pi : \mathcal{E} \rightarrow \mathbb{P}^1$ is elementary and given in detail in [UlmPCMI] so we will not repeat it here. We just note that the fiber of $\mathcal{E}$ over $u = 0$ is a configuration of type I_{2d} and the fibers over the d points where $u^d = 1$ are configuration of type I_2 . The fiber over ∞ is of type I_{2d}^* when d is odd and of type I_{2d} when d is even. All other fibers of π are smooth.

From now on we assume that d has the form $d = p^f + 1$. In particular, d is even.

We label the components of the bad fibers at $u = 0$ and ∞ by the elements of $\mathbb{Z}/2d\mathbb{Z}$ such that the identity component is labeled with 0 and the component meeting the identity component and corresponding to the branch of the reduction of E with tangent line $y = x$ is labeled with 1. The other components are of course labeled cyclically; in particular, the component meeting the identity component and corresponding to the branch of the reduction of E with tangent line $y = -x$ is labelled $2d - 1$.

4.2. Height pairing. The definition of the height pairing is discussed in detail in [UlmPCMI]. We quickly review the construction here. For a point $P \in E(K_d)$, we write $[P]$ for the corresponding curve in $\mathcal{E}$, namely the image of the section $\mathbb{P}^1 \rightarrow \mathcal{E}$ whose generic fiber is P . For each P there is a unique $\mathbb{Q}$ -linear combination D_P of non-identity components of bad fibers of π such that the divisor $[P] - [0] + D_P$ is orthogonal to all components of fibers of π under the intersection pairing on $\mathcal{E}$. To compute D_P , one need only determine the component of the reduction through which $[P]$ passes at each reducible fiber. The height pairing is then defined by

$$(4.2.1) \quad \langle P, Q \rangle = -([P] - [0] + D_P) \cdot ([Q] - [0]) = -([P] - [0]) \cdot ([Q] - [0]) - D_P \cdot [Q].$$

Here the dot signifies the intersection pairing on $\mathcal{E}$. This pairing is known to be symmetric and positive definite modulo torsion (e.g., [CZ79, 1.15]). It makes $E(K_d)$

into a lattice in the real vector space $E(K_d) \otimes_{\mathbb{Z}} \mathbb{R}$. (The Néron-Tate canonical height is $\log q$ times this height. We omit the $\log q$ for simplicity.) The “correction factor” $D_P.[Q]$ is symmetric in P and Q and depends only on the components through which $[P]$ and $[Q]$ pass at each bad fiber. A table of values is given in [CZ79, 1.19].

For any elliptic surface $\pi : \mathcal{E} \rightarrow \mathbb{P}^1$, the self intersection of any section is equal to $-\delta$ where δ is the degree of the invertible sheaf $\omega = (R^1\pi_*\mathcal{O}_{\mathcal{E}})^{-1}$. If $\mathcal{E}$ is non-constant, we also have $\delta = p_g + 1$ where $p_g = \dim H^0(\mathcal{E}, \Omega^2)$ is the geometric genus of $\mathcal{E}$.

4.3. Explicit heights. We apply the above to the elliptic surface $\mathcal{E} \rightarrow \mathbb{P}^1$ associated to E over K_d .

A simple calculation shows that $[P_i].[0] = 0$ for all i and that $[P_i].[P_j] = 0$ if $i \neq j$ in $\mathbb{Z}/d\mathbb{Z}$. The same calculation reveals which components the sections $[P_i]$ pass through: at $u = 0$, all sections $[P_i]$ pass through the component labelled 1; at $u = \infty$, $[P_i]$ passes through the component labelled $d - 1$ if i is even and through the component labelled $d + 1$ if i is odd; at the d points where $u^d = 1$, $[P_i]$ passes through the identity component except at $u = -\zeta_d^{-i}$ where it passes through the non-identity component.

For the elliptic surface associated to the curve E over K_d , we have $\delta = d/2 = (p^f + 1)/2$.

Using the table of corrections appearing in [CZ79] and the calculations above we arrive at the following result.

4.4. Theorem. *Let p be an odd prime number and let E be the elliptic curve over $K = \mathbb{F}_p(t)$ defined by (2.1). Let $d = p^f + 1$ for some $f \geq 0$ and let $K_d = K(\mu_d, u)$ where $u^d = t$. Then the height pairing (4.2.1) of the points P_i ($i = 0, \dots, d - 1$) on E given by (2.2) is*

$$\langle P_i, P_j \rangle = \begin{cases} \frac{(d-1)(d-2)}{2d} & \text{if } i = j \\ \frac{(1-d)}{d} & \text{if } i - j \text{ is even and } \neq 0 \\ 0 & \text{if } i - j \text{ is odd} \end{cases}$$

We write V_d for the subgroup of $E(K_d)$ generated by the P_i and $E(K_d)_{\text{tor}}$. It is easy to see that the lattice V_d modulo torsion is isomorphic to the direct sum of two copies of a scaling of the lattice $A_{d/2-1}^*$ where A_n^* is the dual of the A_n lattice, as in [CS99, 4.6.6]. The points P_i are root vectors; perhaps this explains why it is so easy to write them down.

4.5. Corollary. *The rank of $E(K_d)$ is at least $d - 2$.*

Note that when $d = 2$, the points P_i are independent of p and in fact lift to characteristic zero. The height calculation shows (again) that they are torsion.

For a general $d = p^f + 1$, the group V_d is the direct sum of a torsion group of order 8 and a free group of rank $d - 2$. The determinant of the pairing $\langle, \rangle$ on V_d modulo torsion is

$$2^{4-d}(d-1)^{d-2}d^{-2}.$$

4.6. Regulator and index. We will see below that the index $[E(K_d) : V_d]$ is finite. Assuming this, we have that the regulator of E over $K_d = \mathbb{F}_q(u)$ is

$$R = \frac{2^{4-d}(d-1)^{d-2}d^{-2}}{[E(K_d) : V_d]^2} (\log q)^{d-2}.$$

There is an integrality property of the regulator which affords some control on the index $[E(K_d) : V_d]$ (assuming it is finite). More precisely, for each place v of K_d , let d_v be the discriminant of the intersection pairing on $\mathcal{E}$ restricted to the part of Néron-Severi generated by the non-identity components in the fiber over v . Then it follows from the definition of the height pairing and the integrality of the intersection pairing that the rational number

$$\frac{R/(\log q)^{d-2}}{|E(K_d)_{\text{tor}}|^2} \prod_v d_v$$

is an integer. (This is explained in detail in a more general context in [Gor79, 4.6 and 5.5].) In our setting, the local factors d_v at $u = 0$ and $u = \infty$ are $2d$, at the d places where $u^d = 1$ they are 2, and elsewhere they are 1. It follows that

$$\frac{p^{f(d-2)/2}}{[E(K_d) : V_d]}$$

is an integer. In particular, the index is a power of p .

5. THE TATE-SHAFAREVICH GROUP

We have seen that the curve E has multiplicative reduction at $d + 2$ places of K_d and good reduction elsewhere. It follows that its conductor has degree $d + 2$. By standard results on elliptic curves over function fields, the rank of $E(K_d)$ is bounded above by $d + 2 - 4 = d - 2$ and so the lower bound of Corollary 4.5 is in fact an equality. It also follows that the L function of E over $K_d = \mathbb{F}_q(u)$ has the form

$$L(E/K_d, s) = (1 - q^{1-s})^{d-2}.$$

This follows from the fact that the rank upper bound is an equality; see [UlmPCMI] for more details on this and the rank bound. We could also deduce the L -function from [Ulm07, 4.7] or from the Artin formalism and a root number calculation without knowing anything about points on $E(K_d)$.

Since the order of vanishing of the L -series is equal to the rank (i.e., the basic conjecture of Birch and Swinnerton-Dyer holds), the refined BSD conjecture holds as well and we get an interesting formula for the order of the Tate-Shafarevich group.

Let $\mathbb{III}$ be the Tate-Shafarevich group of E over K_d . The refined conjecture says that

$$L^{(d-2)}(E/K_d, 1) = \frac{|\mathbb{III}| R \tau}{|E(K_d)_{\text{tor}}|^2}.$$

Here R is the regulator calculated in the last section and τ is the Tamagawa number

$$\tau = q^{1-\delta} \prod_v c_v$$

where the product is over the places of K_d and c_v is the order of the group of connected components in the Néron model at v .

Keeping in mind that $K_d = \mathbb{F}_q(u)$ with $\mathbb{F}_q = \mathbb{F}_p(\mu_d)$, we find a beautiful formula for the order of $\mathfrak{III}$:

$$|\mathfrak{III}(E/K_d)| = [E(K_d) : V_d]^2.$$

More generally, if $F = \mathbb{F}_q(u)$ with $u^d = t$, $d = p^f + 1$ and $\mathbb{F}_q$ an extension of $\mathbb{F}_p(\mu_d)$, we find that

$$|\mathfrak{III}(E/F)| = [E(F) : V_d]^2 (q/p^{2f})^{(p^f-1)/2}.$$

Summarizing, we have

5.1. Theorem. *Let p be an odd prime number and let E be the elliptic curve over $K = \mathbb{F}_p(t)$ defined by (2.1). Let $d = p^f + 1$ for some $f \geq 0$ and let $F = \mathbb{F}_q(u)$ where $u^d = t$ and $\mathbb{F}_q$ is a finite extension of $\mathbb{F}_p(\mu_d)$. Then the rank of $E(F)$ is $d-2$ and the Tate-Shafarevich group of E over F is a finite p -group of order at most $q^{(d-2)/2}$. Its order is bounded below by $(q/p^{2f})^{(d-2)/2}$.*

It looks like a very interesting problem to determine exactly the order of $\mathfrak{III}$ and to construct homogeneous spaces representing its elements.

6. EXPLICIT POINTS OVER $\mathbb{F}_p(u)$

Since we have explicit generators for a subgroup of finite index in $E(K_d)$ and the action of $\text{Gal}(K_d/\mathbb{F}_p(u))$ on these generators is evident, it is easy to calculate the rank of E over $\mathbb{F}_q(u)$ for any q . Explicitly, we find:

6.1. Proposition. *Let p be an odd prime number and let E be the elliptic curve over $K = \mathbb{F}_p(t)$ defined by (2.1). Let $d = p^f + 1$ for some $f \geq 0$ and let $F = \mathbb{F}_q(u)$ where $u^d = t$ and $\mathbb{F}_q$ is a finite extension of $\mathbb{F}_p$. Then the rank of $E(F)$ is*

$$\sum_{e|d, e>2} \frac{\phi(e)}{o_q(e)}$$

where the sum is over divisors e of d greater than 2, ϕ is Euler's function, and $o_q(e)$ is the order of q in $(\mathbb{Z}/e\mathbb{Z})^\times$.

Note that when q is a power of p^{2f} , the rank is $p^f - 1$ and when $q = p$, the rank is of the order $p^f/2f$. (See [PS06] for an interesting analysis of the ‘‘average’’ and ‘‘typical’’ values of the rank.)

In particular, E has unbounded rank over the fields $\mathbb{F}_p(t^{1/d})$ for any fixed odd p and varying d . It would be interesting to have explicit generators over these fields.

As a step in this direction, we give explicit generators for a finite index subgroup of $E(\mathbb{F}_p(t^{1/(p+1)}))$. This gives explicit high ranks for sufficiently large p .

6.2. Proposition. *Let p be an odd prime number and let E be the elliptic curve over $K = \mathbb{F}_p(t)$ defined by (2.1). Let $d = p + 1$ and let $F = \mathbb{F}_p(u)$ where $u^d = t$. For each of the $(p-1)/2$ values of $b \in \mathbb{F}_p$ such that $x^2 - bx + 1$ is irreducible in $\mathbb{F}_p[x]$, there is a point $R_b \in E(F)$ of infinite order with x -coordinate*

$$\frac{2u^{p+1} + bu^p + bu + 2 - 2(u^2 + ub + 1)^{d/2}}{b^2 - 4}.$$

These points together with $P_0 = (u, u(u+1)^{d/2})$ and $P_{d/2} = (-u, -u(-u+1)^{d/2})$ generate a subgroup of rank $(p-1)/2$ in $E(F)$.

(Note that the last term in the numerator is $2(u^2 + ub + 1)^{d/2}$ and not $2(u^2 - ub + 1)^{d/2}$.) We could omit P_0 and have the same result. When $p \equiv 1 \pmod{4}$ we could omit both P_0 and $P_{d/2}$.

Proof. Let $\zeta_d \in \mathbb{F}_{p^2}$ be the d -th root of unity fixed above and recall the points P_i defined by (2.2). If the roots of $x^2 - bx + 1$ are ζ^i and $\zeta^{pi} = \zeta^{-i}$, then the point $P_i + P_{-i}$ is defined over $\mathbb{F}_p(t)$. A relatively straightforward computation, which we omit, shows that its x -coordinate is the expression above. That it is of infinite order and that the R_b , P_0 and $P_{d/2}$ generate a group of rank $(p-1)/2$ follows immediately from the height computation Theorem 4.4. $\square$

It does not seem feasible to compute the trace $\sum_{j=0}^{f-1} P_{ip^j}$ by hand when $f > 1$ and so a new idea is needed to find explicit high ranks over $\mathbb{F}_p(u)$ for a fixed p .

7. CONNECTION WITH BERGER'S CONTRUCTION

It is not clear whether the Legendre curve arises via Berger's construction, as in [UlmDPCT, Section 4], but we can make an indirect connection with a conclusion which is just as useful.

Let k be a field of characteristic $\neq 2$ and let E' over $k(t')$ be the elliptic curve studied in Section 7 of [UlmDPCT]:

$$E' : y^2 + xy + t'y = x^3 + t'x^2.$$

We identify the fields $k(t)$ and $k(t')$ by setting $t' = t/16$ and view E' as an elliptic curve over $k(t)$.

7.1. Lemma. *The elliptic curve E' over $k(t)$ is 2-isogenous to the Legendre curve (2.1).*

Proof. This follows from a simple computation. On E' , change coordinates $y \mapsto y - (x + t/16)/2$ to simplify the left hand side, $x \mapsto x - t/16$ to clear the constant term on the right hand side, and $(x, y) \mapsto (x/16, y/64)$ to clear denominators. The result has equation

$$y^2 = x^3 + (4 - 2t)x^2 + t^2x.$$

Dividing by the 2-torsion point $(0, 0)$ yields the curve

$$y^2 = x^3 + (4t - 8)x^2 - 16(t - 1)x = x(x - 4)(x + 4(t - 1)).$$

Finally, the change of coordinates $(x, y) \mapsto (4x + 4, 8y)$ yield the Legendre curve

$$y^2 = x(x + 1)(x + t).$$

$\square$

This lemma allows us to transfer strong results about E' over to E .

7.2. Corollary. *If k is a field of characteristic zero, then the rank of $E(k(t^{1/d}))$ is zero for all d . If k is a finite field of characteristic $p \neq 2$, then the Birch and Swinnerton-Dyer conjecture holds for E over $k(t^{1/d})$ for all d prime to p .*

Proof. The conclusions (rank zero or BSD) both “descend” under finite extensions of k , i.e., if they hold for E over $k'(t^{1/d})$ with k' a finite extension of k , then they hold over $k(t^{1/d})$. So for a fixed d we may assume that k contains a d -th root of 16. In this case, we may take $k(t^{1/d})$ and $k(t'^{1/d}) = k((t/16)^{1/d})$ to be the same field inside an algebraic closure of $k(t)$. The lemma shows that E and E' are 2-isogenous over this field and we proved the claims of the lemma for E' in [UlmDPCT, Section 7]. This shows that these claims also hold for E . $\square$

For k finite and d dividing $p^a + 1$ for some a , we have calculated the rank of $E(k(t^{1/d}))$ now in three different ways: via cohomology in [Ulm07], using Berger's construction and endomorphisms of Jacobians in [UlmDPCT], and "by hand" in this paper. We hope to address the question of the rank for other values of d in a future work.

REFERENCES

- [Ber08] L. Berger, *Towers of surfaces dominated by products of curves and elliptic curves of large rank over function fields*, J. Number Theory **128** (2008), 3013–3030.
- [BDS04] I. I. Bouw, C. Diem, and J. Scholten, *Ordinary elliptic curves of high rank over $\mathbb{F}_p(x)$ with constant j -invariant*, Manuscripta Math. **114** (2004), 487–501.
- [Con09] R. Conceição, *Twists Of elliptic curves over $\mathbb{F}_q(T)$ with a large set of integral points* (2009), Preprint.
- [CS99] J. H. Conway and N. J. A. Sloane, *Sphere packings, lattices and groups*, 3rd ed., Grundlehren der Mathematischen Wissenschaften, vol. 290, Springer-Verlag, New York, 1999.
- [CZ79] D. A. Cox and S. Zucker, *Intersection numbers of sections of elliptic surfaces*, Invent. Math. **53** (1979), 1–44.
- [Dar04] H. Darmon, *Heegner points and elliptic curves of large rank over function fields*, Heegner points and Rankin L -series, Math. Sci. Res. Inst. Publ., vol. 49, Cambridge Univ. Press, Cambridge, 2004, pp. 317–322.
- [Gor79] W. J. Gordon, *Linking the conjectures of Artin-Tate and Birch-Swinnerton-Dyer*, Compositio Math. **38** (1979), 163–199.
- [Mil86] J. S. Milne, *Jacobian varieties*, Arithmetic geometry (Storrs, Conn., 1984), Springer, New York, 1986, pp. 167–212.
- [PS06] C. Pomerance and I. E. Shparlinski, *Rank statistics for a family of elliptic curves over a function field* (2006), To appear in Pure Appl. Math. Q.
- [Shi86] T. Shioda, *An explicit algorithm for computing the Picard number of certain algebraic surfaces*, Amer. J. Math. **108** (1986), 415–432.
- [Shi91] ———, *Mordell-Weil lattices and sphere packings*, Amer. J. Math. **113** (1991), 931–948.
- [Tat66] J. T. Tate, *Endomorphisms of abelian varieties over finite fields*, Invent. Math. **2** (1966), 134–144.
- [TŠ67] J. T. Tate and I. R. Šafarevič, *The rank of elliptic curves*, Dokl. Akad. Nauk SSSR **175** (1967), 770–773 (Russian).
- [Ulm02] D. Ulmer, *Elliptic curves with large rank over function fields*, Ann. of Math. (2) **155** (2002), 295–315.
- [Ulm07] ———, *L -functions with large analytic rank and abelian varieties with large algebraic rank over function fields*, Invent. Math. **167** (2007), 379–408.
- [UlmDPCT] ———, *On Mordell-Weil groups of Jacobians over function fields* (2009), Preprint.
- [UlmPCMI] ———, *Notes on lectures at the 2009 IAS/Park City Mathematics Institute* (2009), In preparation.

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332
E-mail address: `ulmer@math.gatech.edu`